

# VEGFA rs3025039 and biliary atresia susceptibility in Chinese population: a systematic review and meta-analysis

Shiwei He, Yifan Yang, Lingdu Meng, Gong Chen, Yanlei Huang, Zhen Shen, Rui Dong, Shan Zheng

**To cite:** He S, Yang Y, Meng L, et al. VEGFA rs3025039 and biliary atresia susceptibility in Chinese population: a systematic review and meta-analysis. *World Jnl Ped Surg* 2022;5:e000344. doi:10.1136/wjps-2021-000344  
▶ Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/wjps-2021-000344>).

SH, YY and LM contributed equally.

Received 26 July 2021

Accepted 8 November 2021

## ABSTRACT

**Background** Previous studies have suggested an association between vascular endothelial growth factor A (VEGFA) rs3025039 polymorphism and biliary atresia (BA). However, this conclusion is controversial and there is no published pooled evidence of this association.

**Methods** This study was conducted and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses. The protocol was registered with PROSPERO (International Prospective Register of Systematic Reviews). A thorough search was performed on databases including PubMed, Embase, and Chinese Biomedical Database up to August 2020. This study included 846 cases of BA and 2821 controls concerning VEGFA rs3025039 polymorphism. We selected relevant studies based on the following inclusion criteria: (1) the study design was case-control and cohort and (2) the patients carried standard clinical diagnoses of BA, etc. The exclusion criteria were as follows: (1) patients with other related diseases, (2) lack of requisite information and (3) duplicate data. The OR (odds ratio) and the corresponding 95% CI (confidence interval) were calculated to estimate the association.

**Results** This study on VEGFA rs3025039 polymorphism in the Chinese population included 846 cases and 2821 controls. The results showed that there was no significant association between rs3025039 and susceptibility to BA under four genetic models. The results of the subgroup analysis were similar to the overall results.

**Conclusions** This meta-analysis shows that rs3025039 was not associated with susceptibility to BA in the Chinese population. Further validation may entail additional research.

**PROSPERO registration number** CRD42020203812.

## INTRODUCTION

Biliary atresia (BA) is a type of progressive obliterative disorder in neonates that interferes with the function and anatomy of the intrahepatic and extrahepatic bile ducts.<sup>1,2</sup> This destructive inflammatory obliterative cholangiopathy frequently leads to hepatic fibrosis and end-stage liver disease. If untreated, BA with progressive liver cirrhosis is uniformly fatal.<sup>3</sup> The clear etiology of this disorder is not well understood. Genetic

## Key messages

### What is already known about this subject?

- ▶ The etiology of biliary atresia (BA) is still not well understood.
- ▶ Previous studies supported that vascular endothelial growth factor A (VEGFA) plays an important role in the pathogenesis of BA.
- ▶ The association between the single nucleotide polymorphism (SNP) (rs3025039) in the VEGFA gene and the risk of BA is still controversial due to inconsistency among the previous studies.

### What are the new findings?

- ▶ VEGFA rs3025039 polymorphism may not be associated with an elevated risk of BA in Chinese.
- ▶ The association between rs3025039 and risk of BA was not significant in southern Chinese.
- ▶ This meta-analysis did not demonstrate a definite association between rs3025039 and BA in north-western Chinese because only one study was included in this subgroup.

### How might it impact on clinical practice in the foreseeable future?

- ▶ Additional research is needed to obtain a definite conclusion of this association.
- ▶ Future studies are recommended to identify other possible genetic markers.

and immunological factors, infections, and other environmental factors might lead to BA, suggesting that it has a complex etiology.<sup>3,4</sup> Hereditary factors participate in the pathogenesis of BA. Multiple single-nucleotide polymorphisms on the genes, including *ADD3*, *XPNPEP1*, *VEGFA*, and *EFEMP1*, are associated with risk of BA.<sup>4</sup>

The human VEGFA gene is one of the members of the VEGF (vascular endothelial growth factor) family located on chromosome 6p21.3. This gene comprises a type of heparin-binding protein that is in the form of a disulfide-linked homodimer. VEGFA participates in various developmental processes, including endothelial cell proliferation, cell migration,




© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

Department of Pediatric Surgery, Children's Hospital of Fudan University, Shanghai Key Laboratory of Birth Defect, Shanghai, China

## Correspondence to

Dr Shan Zheng; [szheng@shmu.edu.cn](mailto:szheng@shmu.edu.cn)

Dr Rui Dong; [rdong@fudan.edu.cn](mailto:rdong@fudan.edu.cn)



**Figure 1** Flow chart of studies considered for inclusion. BA, biliary atresia.

and apoptosis.<sup>5-7</sup> It may also participate in the pathogenesis of BA because it can function as a proinflammatory cytokine.<sup>8</sup> Polymorphisms of this functional gene may affect expression regulation, leading to various incidences and severities of disease.<sup>9</sup> Thus, clarifying the effects of alterations within the *VEGFA* gene may provide markers for diagnosis and treatment to reverse progression of BA.

Previous studies have explored the correlation between rs3025039 within the *VEGFA* gene and susceptibility to BA; however, the results remain controversial due to the lack of consistency among the studies. In China, rs3025039 has been found to be associated with susceptibility to BA.<sup>10-11</sup> Interestingly, a recent publication reported no significant correlation between rs3025039 and BA in the southern Chinese population.<sup>12</sup>

This discrepancy needs further verification by increasing the sample size. In the mean time, a meta-analysis is a suitable method to summarize previous genetic association studies (GAS) and to draw relatively reliable conclusions.<sup>13-14</sup> Therefore, the present meta-analysis may provide evidence regarding the association of *VEGFA* rs3025039 polymorphism with susceptibility to BA.

## METHODS

This study protocol was registered with an international registration platform of systematic review, PROSPERO (International Prospective Register of Systematic Reviews). We conducted and reported the study according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement<sup>15</sup> (see online supplemental table S1).

### Patient and public involvement

Patients and the public were not involved in this study.

### Data sources and searches

PubMed, Embase, and the Chinese Biomedical Database (CBD) were searched from inception until August 17, 2020 (see online supplemental material for full details on search strategy). We also considered references in the included studies and related reviews. We did not impose limitations on the language of papers, time period of follow-up, and published state. We reran the same searches before the final analyses and retrieved additional studies for inclusion.

### Eligibility criteria

We established the inclusion and exclusion criteria based on discussion studies. We selected relevant studies based

**Table 1** Characteristics of the studies included in this meta-analysis of VEGFA rs3025039 polymorphism

Study	Population	Area	Race	Sample size				Genotype in				Minor allele frequency in controls (%)	HWE p value in controls	Type of study	Genotyping method	Quality score
				Cases	Controls	CC	CT	TT	CC	CT	TT					
Lee et al <sup>10</sup>	Chinese	Southern	Asian	45	160	37	8	0	91	62	7	23.8	>0.05	Hospital-based	TaqMan	6
Liu et al <sup>11</sup>	Chinese	Northwestern	Asian	311	1205	220	86	5	620	355	50	22.2	0.929	Hospital-based	MassARRAY	6
Liu et al <sup>12</sup>	Chinese	Southern	Asian	490*	1456*	332	142	16	1002	411	43	17	0.874	Hospital-based	MassARRAY	9

\*The number of cases and controls was calculated from the publication. HWE, Hardy-Weinberg equilibrium; VEGFA, vascular endothelial growth factor A.

on the following inclusion criteria: (1) the study design was case-control and cohort; (2) the patients carried a standard clinical diagnosis of BA; (3) the studies explored the target association; and (4) the authors presented enough data on genotype distribution. The criteria used to exclude studies were as follows: (1) patients with other related diseases; (2) lack of requisite information; and (3) duplicate data.

### Study selection

At the first stage, duplicates from three electronic databases were screened and removed independently by three reviewers (SH, YY, and LM). At the second stage, the title and abstract of each of the remaining studies were reviewed independently by the same reviewers (SH, YY, and LM) to select eligible studies. At the final stage, the same reviewers (SH, YY, and LM) independently retrieved and assessed the potentially eligible full text of the remaining publications. Any discrepancy with regard to eligibility of articles was discussed among the three reviewers in consultation with a third reviewer (RD or SZ).

### Data extraction

All data were recorded independently by three reviewers (SH, YY, and LM) in accordance with a record form, with regard to (1) study characteristics (author information, publication time, sample size, country, and ethnic origin); and (2) genotype data (number with different genotypes, minor allele frequency, results of the Hardy-Weinberg equilibrium (HWE) of the control group, and genotyping methods). All related data were found in the original studies so we did not need to contact the study authors to request for missing data.

### Quality score assessment

Three reviewers (SH, YY, and LM) independently appraised the quality of the GAS using a checklist revised from previous studies,<sup>16 17</sup> which was done on the basis of genetic factors and epidemiological requirements. The checklist covered essential items aimed at the quality of the GAS, including representativeness and ascertainment of study subjects, genotyping, HWE, and association analysis.<sup>16 17</sup> The total score ranged from 0 (worst) to 13 (best). Detailed information is shown in online supplemental table S2.

### Statistical analysis

In a GAS, research candidates should be categorized into three groups (BB, Bb, and bb) and usually B is used as the susceptibility allele.<sup>14 18 19</sup> Previous studies have suggested that the C allele increased BA susceptibility<sup>11</sup>; therefore, we estimated the association between rs3025039 and susceptibility to BA using four different genetic models, namely a per-allele model (C vs T), a homozygous model (CC vs TT), a dominant model (CC+CT vs TT), and a recessive model (CC vs CT+TT).<sup>19</sup> We measured the effects using OR (odds ratio) and 95% CI (confidence interval) using a fixed-effect or a random-effect model.<sup>20</sup> We assessed heterogeneity using the Cochrane Q statistic and the inconsistency index ( $I^2$ ).<sup>21</sup>  $I^2 > 50\%$  or p value  $< 0.1$  indicated substantial heterogeneity.

**Table 2** Main results of the meta-analysis

Studies (n)	Comparison model	Test of association			Test of heterogeneity			P value for publication bias	
		OR	95% CI	P value	Q	P value	I <sup>2</sup> (%)	Begg's test	Egger's test
3	C vs T	1.50	0.90 to 2.49	0.121	16.52	0.001	87.9	0.602	0.422
3	CC vs TT	2.02	0.58 to 6.99	0.269	7.49	0.024	73.3		
3	CT+CC vs TT	1.78	0.60 to 5.32	0.299	2.96	0.052	66.2		
3	CC vs CT+TT	1.53	0.89 to 2.63	0.124	5.92	0.001	86.3		

CI, confidence interval; OR, odd ratio.

We conducted subgroup analyses according to the several study subject areas. We performed a sensitivity analysis by excluding every publication individually to evaluate the reliability and stability of the overall OR. We appraised publication bias using Egger's test and Begg's test<sup>22</sup> and visual inspection of funnel plots.<sup>23</sup> The meta package (V.4.9.7) in R software (V.3.6.1) was used to complete all analyses. In addition to heterogeneity,  $p < 0.05$  (two-tailed) indicated significance.

## RESULTS

### Search findings

We conducted the search process and reported the findings according to the PRISMA statement<sup>15</sup> (see figure 1). We identified 41 papers after an initial search. At the first stage, we removed 10 duplicate articles, leaving 31 articles of potential relevance. At the second stage, we excluded 5 papers that did not involve patients with BA, 5 studies on animal experiments, 7 traditional reviews, and 10 studies that were not about polymorphisms. At the final stage, we excluded one paper due to insufficient information for inclusion. Finally, we included three articles<sup>10–12</sup> for data extraction and meta-analysis.

### Study characteristics

Table 1 shows the main data of the three studies (846 cases and 2821 controls). All of the included studies were conducted in the Chinese population. The genotype distributions of the controls in the three studies<sup>10–12</sup> were consistent with HWE, but one did not provide exact data.<sup>12</sup> Of all studies, two<sup>10,12</sup> involved southern Chinese individuals, and one study<sup>11</sup> involved northwestern Chinese individuals. All studies<sup>10–12</sup> used a case–control design. One study<sup>12</sup> was a case–control cohort study, and another study<sup>11</sup> was a

case–control study but did not report the specific type. All studies used hospital-based controls. The total scores for the three studies ranged from 6 to 9 (see table 1 and online supplemental table S3).

### Heterogeneity test


There was a significant between-study heterogeneity in four genetic models of rs3025039 polymorphism (I<sup>2</sup> range: 66.2%–87.9%,  $p = 0.0003$ – $0.0518$ ; see table 2). Therefore, we used a random-effects model to combine the associations between rs3025039 polymorphism and risk of BA.

### Association between rs3025039 and risk of BA

Three studies, including 846 cases and 2821 controls, examined the association between rs3025039 and susceptibility to BA. There was no statistically significant association in any of the genetic models (C vs T: OR=1.50, 95% CI 0.90 to 2.49; CC vs TT: OR=2.02, 95% CI 0.58 to 6.99; CT+CC vs TT: OR=1.78, 95% CI 0.60 to 5.32; CC vs CT+TT: OR=1.53, 95% CI 0.89 to 2.63). Table 2 and figure 2 display the distribution of the rs3025039 genotypes and alleles.

### Subgroup analysis

We performed a subgroup analysis by area. Table 3 details the results. Southern Chinese individuals were the subjects in two studies, while one study involved northwestern Chinese individuals. The association between rs3025039 and susceptibility to BA was not significant in southern Chinese patients (C vs T: OR=1.64, 95% CI 0.50 to 5.37; CC vs TT: OR=1.40, 95% CI 0.27 to 7.14; CT+CC vs TT: OR=1.06, 95% CI 0.40 to 2.80; CC vs CT+TT: OR=1.72, 95% CI 0.48 to 6.14). For the northwestern Chinese subgroup, we could not perform a meta-analysis because there was only one study.



**Figure 2** Forest plot of per-allele model of the association between VEGFA rs3025039 polymorphism and biliary atresia. VEGFA, vascular endothelial growth factor A.

**Table 3** Results of the subgroup analysis

Area	Studies (n)	Comparison model	OR	95% CI
Southern	2	C vs T	1.64	0.50 to 5.37
	2	CC vs TT	1.40	0.27 to 7.14
	2	CT+CC vs TT	1.06	0.40 to 2.80
	2	CC vs CT+TT	1.72	0.48 to 6.14

CI, confidence interval; OR, odd ratio.

### Sensitivity analysis

A sensitivity analysis was performed to evaluate the effects of individual studies on the pooled OR. The pooled OR with 95% CI changed after omitting Liu *et al*<sup>12</sup> (OR=2.03, 95% CI 1.03 to 3.98), suggesting that this study was a source of heterogeneity ( $I^2$  decreased from 87.9% to 66.9%) (figure 3).

### Publication bias

Both tests supported the absence of publication bias (p value for Begg's test=0.602, p value for Egger's test=0.422; table 2). However, the shape of the funnel plot was asymmetric (figure 4). Because a funnel plot requires at least five studies and because we considered only three studies, there was a possibility that there was a possibility for the funnel plot to show asymmetry. Therefore, we considered only the results of the Begg's and Egger's tests.

### DISCUSSION


We conducted a systematic review of the association between *VEGFA* rs3025039 polymorphism and susceptibility to BA. Lee *et al*<sup>10</sup> were the first to identify *VEGFA* as a susceptibility locus for BA in Chinese patients. This result has biological plausibility; *VEGFA* is a mediator of the pathogenesis of BA. As a proinflammatory cytokine, *VEGFA* participates in various processes including cell proliferation, cell migration, and apoptosis, all of which occur in cell-mediated immun-inflammatory diseases, such as BA.<sup>5 6 8</sup> There are several independent verifications of this association, but the conclusions remain controversial because the results of different studies in China were conflicting.<sup>11 12</sup>


We found a difference between cases and controls in comparisons of all genotypes of rs3025039, suggesting that rs3025039 polymorphism may not correlate with susceptibility to BA. A subgroup analysis by area further suggested no significant association between rs3025039 and susceptibility to BA. Given that only one study included the subgroup

of northwestern Chinese, our study did not demonstrate a solid correlation between rs3025039 and BA in northwestern Chinese patients.

The absence of significant findings may stem from the fact that we included only three original studies in this study. In addition, as the sample size increased, the association between rs3025039 and susceptibility to BA decreased in the three studies. We observed statistical significance in only two studies with small numbers of participants. Therefore, the statistical power was relatively low, suggesting that the association might be spurious. For the first reason, more original studies regarding the association between rs3025039 and susceptibility to BA are necessary to generate accurate results. However, for the second reason, the effect may be relatively small or not at all, and explicit exploration of this association may result in an unnecessary study. Therefore, it may be more cost-effective to devote more resources to explore other potential biomarkers.

Several limitations need to be acknowledged when considering the results of our study. First, the quantity and sample size of the included studies were insufficient to obtain high power for making a confirmatory conclusion, even though we undertook a comprehensive literature search. Second, all study subjects came from China only; therefore, we could not avoid potential selection bias. Third, residual confounders were possible because BA is a multifactorial and complicated disease, involving gene–environment and gene–gene interactions. We could not detect these effects due to limited information. Fourth, there was a lack of sufficient data; therefore, we did not conduct a subgroup analysis of familial and other BA types. Finally, there was significant heterogeneity that might distort the results. Several aspects may cause heterogeneity, including differences in experimental methods across studies. Therefore, readers should interpret our results with caution.


**Figure 3** Forest plot of per-allele model for sensitivity analysis.



**Figure 4** Begg's funnel plot of publication bias.

Despite the limitations, we believe that the present study provides useful evidence regarding the role of *VEGFA* rs3025039 polymorphism in BA. The sample size of each study included in this study was insufficient to make a definite conclusion regarding the association between rs3025039 polymorphism and susceptibility to BA; however, the pooled OR calculated from the three studies significantly increased the statistical power. It is worth noting that the sample size increased from 205 subjects in the first study to 3667 subjects in our study. Reaching a relative sufficient statistical power is crucial in GAS. Furthermore, no significant publication bias was found in this study and the result was relatively stable in the subgroup analysis.

In conclusion, *VEGFA* rs3025039 polymorphism might not be associated with an elevated risk of BA in Chinese population. Obtaining a definitive conclusion regarding this association may entail additional research. Future studies are recommended to identify other possible genetic markers.

**Contributors** SH contributed to conceptualization, data curation, formal analysis, project administration, resources, investigation, methodology, software, visualization, writing - original draft. YY contributed to data curation, formal analysis, resources, funding acquisition, investigation, methodology, software, writing - review and editing. LM contributed to data curation, formal analysis, resources, investigation, methodology, visualization, writing - review and editing. GC contributed to resources, writing - review and editing. YH contributed to supervision, writing - review and editing. ZS contributed to supervision, resources. RD contributed to conceptualization, project administration, funding acquisition, supervision, validation, writing - review and editing. SZ contributed to project administration, funding acquisition, supervision, validation, writing - review and editing. All authors reviewed the results and approved the final version of the manuscript.

**Funding** This study received financial support from the Clinical Research Plan of SHDC (no SHDC2020CR2009A), Shanghai Municipal Key Clinical Specialty (no shslczdzk05703), National Natural Science Foundation of China (no 81770519, no 81771633, no 81873545, and no 81974059), Science Foundation of Shanghai (no 18411969100 and no 19ZR1406600), and Children's National Medical Center (no EK1125180104, no EKYY20180204, no EK112520180211, and no EK112520180310).

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** This study is a systematic review and meta-analysis and the data from this study were extracted from the original studies. Moreover, there are no

patients or animals involved in this study. Thus, this study does not require ethics approval.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open access repository. As a meta-analysis, all of the data in this study can be found in table 1 and were extracted from the original studies.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

## REFERENCES

- Asai A, Miethke A, Bezerra JA. Pathogenesis of biliary atresia: defining biology to understand clinical phenotypes. *Nat Rev Gastroenterol Hepatol* 2015;12:342–52.
- Hartley JL, Davenport M, Kelly DA. Biliary atresia. *The Lancet* 2009;374:1704–13.
- Verkade HJ, Bezerra JA, Davenport M, *et al*. Biliary atresia and other cholestatic childhood diseases: advances and future challenges. *J Hepatol* 2016;65:631–42.
- Hartley JL, Davenport M, Kelly DA. Biliary atresia. *Lancet* 2009;374:1704–13.
- Detmar M, Brown LF, Schön MP, *et al*. Increased microvascular density and enhanced leukocyte rolling and adhesion in the skin of VEGF transgenic mice. *J Invest Dermatol* 1998;111:1–6.
- Mor F, Quintana FJ, Cohen IR. Angiogenesis-inflammation cross-talk: vascular endothelial growth factor is secreted by activated T cells and induces Th1 polarization. *J Immunol* 2004;172:4618–23.
- Cohen T, Nahari D, Cerem LW, *et al*. Interleukin 6 induces the expression of vascular endothelial growth factor. *J Biol Chem* 1996;271:736–41.
- Enjoji M, Nakamura M, Yamaguchi K, *et al*. Clinical significance of serum levels of vascular endothelial growth factor and its receptor in biliary disease and carcinoma. *World J Gastroenterol* 2005;11:1167–71.
- Katsonis P, Koire A, Wilson SJ, *et al*. Single nucleotide variations: biological impact and theoretical interpretation. *Protein Sci* 2014;23:1650–66.
- Lee H-C, Chang T-Y, Yeung C-Y, *et al*. Genetic variation in the vascular endothelial growth factor gene is associated with biliary atresia. *J Clin Gastroenterol* 2010;44:135–9.
- Liu B, Wei J, Li M, *et al*. Association of common genetic variants in *VEGFA* with biliary atresia susceptibility in northwestern Han Chinese. *Gene* 2017;628:87–92.
- Liu F, Zeng J, Zhu D, *et al*. Association of polymorphism in the *VEGFA* gene 3'-UTR +936T/C with susceptibility to biliary atresia in a southern Chinese Han population. *J Clin Lab Anal* 2018;32:e22342–5.
- Munafo MR, Flint J. Meta-analysis of genetic association studies. *Trends Genet* 2004;20:439–44.
- Nakaoka H, Inoue I. Meta-analysis of genetic association studies: methodologies, between-study heterogeneity and winner's curse. *J Hum Genet* 2009;54:615–23.
- Moher D, Liberati A, Tetzlaff J, *et al*. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
- Jiang M, Li C, Cao G, *et al*. Effects of *NRG1* polymorphisms on Hirschsprung's disease susceptibility: a meta-analysis. *Sci Rep* 2017;7:9913.
- Thakkestian A, McEvoy M, Minelli C, *et al*. Systematic review and meta-analysis of the association between {beta}2-adrenoceptor polymorphisms and asthma: a HuGE review. *Am J Epidemiol* 2005;162:201–11.
- Lee YH. Meta-Analysis of genetic association studies. *Ann Lab Med* 2015;35:283–7.
- Attia J, Thakkestian A, D'Este C. Meta-analyses of molecular association studies: methodologic lessons for genetic epidemiology. *J Clin Epidemiol* 2003;56:297–303.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- Hayashino Y, Noguchi Y, Fukui T. Systematic evaluation and comparison of statistical tests for publication bias. *J Epidemiol* 2005;15:235–43.
- Egger M, Davey Smith G, Schneider M, *et al*. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.

PubMed (1900/00/00-2020/08/18)

(((((Biliary atresia[MeSH Terms]) OR (((((((((Biliary atresia[Title/Abstract]) OR (Atresia, Biliary[Title/Abstract])) OR (Intrahepatic Biliary Atresia[Title/Abstract])) OR (Atresia, Intrahepatic Biliary[Title/Abstract])) OR (Biliary Atresia, Intrahepatic[Title/Abstract])) OR (Biliary Atresia, Extrahepatic[Title/Abstract])) OR (Atresia, Extrahepatic Biliary[Title/Abstract])) OR (Extrahepatic Biliary Atresia[Title/Abstract])) OR (Idiopathic Extrahepatic Biliary Atresia[Title/Abstract])) OR (Familial Extrahepatic Biliary Atresia[Title/Abstract]))) AND (((((((((((Vascular Endothelial Growth Factor-A[Title/Abstract]) OR (VEGF-A[Title/Abstract])) OR (VEGFA[Title/Abstract])) OR (Vasculotropin[Title/Abstract])) OR (VEGF[Title/Abstract])) OR (Vascular Endothelial Growth Factor[Title/Abstract])) OR (Vascular Permeability Factor[Title/Abstract])) OR (Permeability Factor, Vascular[Title/Abstract])) OR (Glioma-Derived Vascular Endothelial Cell Growth Factor[Title/Abstract])) OR (Glioma Derived Vascular Endothelial Cell Growth Factor[Title/Abstract])) OR (GD-VEGF[Title/Abstract])) OR ((Vascular Endothelial Growth Factor A[MeSH Terms]) OR (Vascular Endothelial Growth Factors[MeSH Terms]))) AND (("1900/01/01"[Date - Publication] : "2020/08/18"[Date - Publication]))

Embase (1900/00/00-2020/08/18)

No.	Query	Results	Date
#30	#12 AND #27 AND [1-1-1900]/sd NOT [18-8-2020]/sd	28	20-Aug-20
#29	#12 AND #27 AND [1-1-1900]/sd NOT [19-8-2020]/sd	28	20-Aug-20
#28	#12 AND #27	28	20-Aug-20
#27	#13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26	151830	20-Aug-20
#26	vascular endothelial growth factors':ti,ab	1250	20-Aug-20
#25	vascular endothelial growth factor a':ti,ab	4416	20-Aug-20
#24	gd-vegf':ti,ab	2	20-Aug-20
#23	glioma derived vascular endothelial cell growth factor':ti,ab	1	20-Aug-20
#22	glioma-derived vascular endothelial cell growth factor':ti,ab	1	20-Aug-20
#21	permeability factor, vascular':ti,ab	123	20-Aug-20
#20	vascular permeability factor':ti,ab	687	20-Aug-20
#19	vascular endothelial growth factor':ti,ab	74734	20-Aug-20
#18	vegf':ti,ab	98548	20-Aug-20
#17	vasculotropin':ti,ab	13	20-Aug-20
#16	vegfa':ti,ab	12762	20-Aug-20
#15	vegf-a':ti,ab	9885	20-Aug-20

#14	vascular endothelial growth factor-a':ti,ab	4416	20-Aug-20
#13	vasculotropin'/exp	109406	20-Aug-20
#12	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11	8458	20-Aug-20
#11	familial extrahepatic biliary atresia':ti,ab	5	20-Aug-20
#10	idiopathic extrahepatic biliary atresia':ti,ab	1	20-Aug-20
#9	extrahepatic biliary atresia':ti,ab	615	20-Aug-20
#8	atresia, extrahepatic biliary':ti,ab	5	20-Aug-20
#7	biliary atresia, extrahepatic':ti,ab	1	20-Aug-20
#6	biliary atresia, intrahepatic':ti,ab	7	20-Aug-20
#5	atresia, intrahepatic biliary':ti,ab	0	20-Aug-20
#4	intrahepatic biliary atresia':ti,ab	34	20-Aug-20
#3	atresia, biliary':ti,ab	27	20-Aug-20
#2	biliary atresia':ti,ab	6581	20-Aug-20
#1	bile duct atresia'/exp	7634	20-Aug-20

## CBM (1900-2020)

("胆道闭锁"[摘要]) AND (("血管内皮生长因子 A"[摘要] OR "血管内皮生长因子"[摘要] OR "VEGFA"[摘要] OR "VEGF"[摘要] OR "Vascular Endothelial Growth Factor A"[摘要] OR "Vascular Endothelial Growth Factors"[摘要] OR "Vascular Endothelial Growth Factor"[摘要]) AND 1900-2020[日期])



**Table S1: The checklist of Preferred Reporting Items for Systematic Reviews and Meta-Analyses**

Section/topic	#	Checklist item	Reported on page #
<b>TITLE</b>			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Page 1
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	Page 1
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known.	Page 2
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	Page 2
<b>METHODS</b>			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	Page 3
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	Page 3
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	Page 3
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Page 3
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	Page 3
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	Page 3
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	Page 3
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	Page 3
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	Page 3
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., $I^2$ ) for each meta-analysis.	Page 3-4

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	Page 3
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	Page 4
<b>RESULTS</b>			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Page 5
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	Page 5
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Page 5
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Page 5
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	Page 5
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Page 5
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	Page 5
<b>DISCUSSION</b>			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	Page 6
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	Page 6
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	Page 6
<b>FUNDING</b>			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	Title page

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: [www.prisma-statement.org](http://www.prisma-statement.org).

**Table S2: Scale for quality assessment of selected study.**

<b>Criteria</b>	<b>Score</b>
<b>Representativeness of cases</b>	
Consecutive/randomly selected from case population with clearly defined sampling frame	2
Consecutive/randomly selected from case population without clearly defined sampling frame or with extensive inclusion/exclusion criteria	1
No method of selection described	0
<b>Representativeness of controls</b>	
Controls were consecutive/randomly drawn from the same sampling frame (ward/community) as cases	2
Controls were consecutive/randomly drawn from a different sampling frame as cases	1
Not described	0
<b>Ascertainment of BA</b>	
Clearly described objective criteria for diagnosis of BA	2
Diagnosis of BA by patient self-report or by patient history	1
Not described	0
<b>Ascertainment of controls</b>	
Controls were ruled out BA. i.e., perform laparoscopic cholangiography, biopsy of the liver and extrahepatic biliary tree or clinical followed up	2
Controls were subjects who did not report BA; no objective testing	1
Not described	0
<b>Genotyping examination</b>	
Genotyping done under “blinded” condition	1
Un-blinded or not mentioned	0
<b>Hardy-Weinberg equilibrium</b>	
Hardy-Weinberg equilibrium in control group	2
Hardy-Weinberg disequilibrium in control group	1
No checking for Hardy-Weinberg equilibrium	0
<b>Association assessment</b>	
Assess association between genotypes and BA with appropriate statistics and adjustment for confounders	2
Assess association between genotypes and BA with appropriate statistics without adjustment for confounders	1
Inappropriate statistics used	0

Abbreviation: BA, biliary atresia.

**Table S3: Quality assessment of included studies.**

<b>Study</b>	<b>Representativeness of cases</b>	<b>Representativeness of controls</b>	<b>Ascertainment of BA</b>	<b>Ascertainment of controls</b>	<b>Genotyping examination</b>	<b>Hardy-Weinberg equilibrium</b>	<b>Association assessment</b>	<b>Total score</b>
Lee, H. C. et al 2010	0	0	2	1	0	2	1	6
Liu, B. et al 2017	0	0	2	1	0	2	1	6
Liu, F. et al 2018	0	2	2	1	0	2	2	9

Abbreviation: BA, biliary atresia.